

24. What is the Origin of the Hydrous Micas of Fennoscandia?

By


I. Th. Rosenqvist

When, in 1953, ERIK NORIN described the formation of illitic mica minerals in the sediments in the central Tyrrhenian Sea, he was fully aware of the fact that such minerals, which form the main mass of the active minerals in the Scandinavian clays, can be formed from non-mica minerals within a, geologically speaking, very brief space of time. The great similarity between the illitic micas and the rock-forming micas has otherwise in Scandinavia led to the opinion that the illitic minerals in our Fennoscandian Pleistocene clays have essentially been formed by a slight chemical transformation of finely ground mica freed from the solid rocks by mechanical erosion mainly during the Ice Age. It has, however, always been difficult to explain the great quantities of hydrous mica minerals in the thick Scandinavian clay deposits exclusively by the separation of rock-forming mica from disintegrated rock material. A reasonably well-founded calculation of the quantities of clays below the surface of the Baltic Sea and the North Sea, together with the quantities that are found upon the terrestrial surface of Scandinavia, shows that we have to do with quantities of hydrous mica of an order of magnitude of 10^{12} m³. If, at the same time, we take into consideration the relatively sparse content of micas in gneisses, granites, and gabbros compared with the content in these rocks of quartz and feldspar, we are driven to the conclusion that the purely mechanical disintegration required to supply this entire mass of hydrous mica must simultaneously have resulted in immense quantities of sand and silt. Altogether, such a calculation will be uncertain, in part very uncertain, but it seems nevertheless as if the known quantities of sand could hardly be brought into agreement with the great masses of hydrous mica minerals, if these latter are supposed to be derived exclusively from the disintegration of crystalline rocks.

The problem loses some of its difficulties if we admit the possibility that the clays have been formed essentially by the disintegration of unmetamorphosed or little metamorphosed Cambro-Silurian clay sediments. But this train of thought also leads to geological difficulties, since the areas of low metamorphic or unmetamorphosed Cambro-Silurian rocks that now are available in Scandinavia constitute a comparatively small portion of the Scandinavian rock-floor. Their erosion in the course of the Last Glaciation has not been particularly

mica can drop below 30% of that in the rock-forming mica. Such far-reaching leaching out has in general been assumed as leading to "mixed-layer" minerals between montmorillonite and illite. It seems, however, as if under specific conditions a fairly thorough removal of K^+ and replacement by H_3O^+ with retention of the mica structure could be obtained.

Even if from the formal point of view the change from mica to montmorillonite seems to consist in a replacement of aluminium by silicon, while at the same time the univalent cations are removed, this phenomenon is nevertheless a typical process of oxidization, since it implies a change from a trivalent to a tetravalent ion. In processes that take place according to the following equation, the access of oxygen constitutes a controlling component:



It is therefore imaginable that the minerals dealt with by LINDQVIST have been formed under less oxidizing conditions than in the uppermost crust of the Norwegian soils examined by us, where we believe we have demonstrated the occurrence of a certain amount of montmorillonite of secondary type.

Recently ODD GJEMS (1960) at the Soil Laboratory of the Agricultural University of Norway has found montmorillonite as dominating clay mineral in fractions finer than 2μ in the A₂ horizon in a number of podsol profiles in Fennoscandian clays even where the soil profile is no older than 300 years. Under the conditions that led to the development of podsol profiles the development of montmorillonite is thus definitely established in the Fennoscandian climate. It is, on the other hand, well known that montmorillonite minerals with access to alkalies, especially potassium, can in their turn easily be transformed into mixed-layer minerals, as shown, e.g., by ANNE MARIE BYSTRÖM (1957) for the bentonites in Mount Kinnekulle in South-western Sweden. It has therefore to be considered a very simple chemical process that ultrafine montmorillonite which has been transported by fresh water into the sea can under reducing conditions easily be recrystallized with the formation of mixed-layer minerals, or hydrous mica.

Since each link in such a chain of development is known from nature, it seems reasonable to assume that also a greater or smaller part of our clays has originated in this way. We are still in need of quantitative investigations, and it might be a worth-while task for future sedimentologists to study the complex of problems linked with the mineral weathering of the temperate climate and the sedimentation along our coasts, under oxidizing as well as under reducing conditions. The work inaugurated by ERIK NORIN demands a continuation.

References

- BYSTRÖM, ANNE MARIE, 1957: The clay minerals in the Ordovician bentonite beds in Billingen, South-west Sweden. *Geol. Fören. Förh.* Bd 79, H. 1. Stockholm.
- GJEMS, ODD, 1960: Clay minerals in podsol profiles in Fennoscandia. *Clay Min. Bull.* Vol. 4, No. 24. Galashiels.
- LINDQVIST, BENGT, 1959: The clay minerals of the Yühuat'ai vermiculated clay at Kinkian in the Lower Yangtze Valley, China. *Bull. Geol. Inst. Univ. Uppsala*, Vol. XXXVIII. Uppsala.
- MOUM, J. and ROSENQVIST, I. TH., 1955: Kjemisk bergartsforvitring belyst ved en del leirprofiler. *Norsk Geol. Tidsskr.*, Nr. 34. Bergen.
- 1957: On the weathering of young marine clay. *Norwegian contrib. to the 4th Int. Conf. on Soil Mechanics and Foundation Engineering*, No. 1a/18, pp. 5-7. London.
- NORIN, ERIC, 1953: Occurrence of authigenous illitic mica in the sediments of the central Tyrrhenian Sea. *Bull. Geol. Inst. Univ. Uppsala*, Vol. XXXIV. Uppsala.
- ROSENQVIST, I. TH., 1955 a: Cation exchange capacity of soil determined by calcium-45. *Norges Geotekn. Inst. Publ.*, nr. 8. Oslo.
- 1955 b: Bidrag til Østlandsleirenes petrografi. *Norsk Geol. Tidsskr.*, nr. 35. Bergen.
- SKEMPTON, A. W., 1953: The colloidal "activity" of clays. *3rd Int. Conf. on Soil Mechanics and Foundation Engineering*, Vol. 1, pp. 57-61. Zürich.